



Laparoscopic surgery for complex and recurrent Crohn's disease

Sevim, Yusuf; Akyol, Cihangir; Aytac, Erman; Baca, Bilgi; Bulut, Orhan; Remzi, Feza H

Published in:
World Journal of Gastrointestinal Endoscopy

DOI:
[10.4253/wjge.v9.i4.149](https://doi.org/10.4253/wjge.v9.i4.149)

Publication date:
2017

Document version
Publisher's PDF, also known as Version of record

Document license:
[CC BY-NC](#)

Citation for published version (APA):
Sevim, Y., Akyol, C., Aytac, E., Baca, B., Bulut, O., & Remzi, F. H. (2017). Laparoscopic surgery for complex and recurrent Crohn's disease. *World Journal of Gastrointestinal Endoscopy*, 9(4), 149-152.
<https://doi.org/10.4253/wjge.v9.i4.149>

World Journal of *Gastrointestinal Endoscopy*

World J Gastrointest Endosc 2017 April 16; 9(4): 149-203



EDITORIAL

- 149 Laparoscopic surgery for complex and recurrent Crohn's disease
Sevim Y, Akyol C, Aytac E, Baca B, Bulut O, Remzi FH

MINIREVIEWS

- 153 Non-functioning pancreatic neuroendocrine tumors: Surgery or observation?
Bar-Moshe Y, Mazeh H, Grozinsky-Glasberg S

ORIGINAL ARTICLE

Case Control Study

- 162 Comparative study of outcomes following laparoscopic Roux-en-Y gastric bypass and sleeve gastrectomy in morbidly obese patients: A case control study
Garg H, Priyadarshini P, Aggarwal S, Agarwal S, Chaudhary R

Retrospective Cohort Study

- 171 Does serotonin reuptake inhibitor therapy increase the risk of post-sphincterotomy bleeding in patients undergoing endoscopic retrograde cholangio-pancreatography?
Yadav D, Vargo J, Lopez R, Chahal P

Retrospective Study

- 177 Does deep sedation with propofol affect adenoma detection rates in average risk screening colonoscopy exams?
Thirumurthi S, Raju GS, Pande M, Ruiz J, Carlson R, Hagan KB, Lee JH, Ross WA

- 183 Endoscopic balloon catheter dilatation *via* retrograde or static technique is safe and effective for cricopharyngeal dysfunction
Chandrasekhara V, Koh J, Lattimer L, Dunbar KB, Ravich WJ, Clarke JO

- 189 Analysis of the risk factors for severity in post endoscopic retrograde cholangiopancreatography pancreatitis: The indication of prophylactic treatments
Matsubara H, Urano F, Kinoshita Y, Okamura S, Kawashima H, Goto H, Hirooka Y

Observational Study

- 196 Endoscopic assessment and management of sporadic duodenal adenomas: The results of single centre multidisciplinary management
Rajkomar K, Kweon M, Khan I, Frankish P, Rodgers M, Koea JB

Contents

World Journal of Gastrointestinal Endoscopy
Volume 9 Number 4 April 16, 2017

ABOUT COVER

Editorial Board Member of *World Journal of Gastrointestinal Endoscopy*, Andreas Sieg, MD, PhD, Assistant Professor, Department of Gastroenterology, Practice of Gastroenterology, 69115 Heidelberg, Germany

AIM AND SCOPE

World Journal of Gastrointestinal Endoscopy (*World J Gastrointest Endosc*, *WJGE*, online ISSN 1948-5190, DOI: 10.4253) is a peer-reviewed open access (OA) academic journal that aims to guide clinical practice and improve diagnostic and therapeutic skills of clinicians.

WJGE covers topics concerning gastroscopy, intestinal endoscopy, colonoscopy, capsule endoscopy, laparoscopy, interventional diagnosis and therapy, as well as advances in technology. Emphasis is placed on the clinical practice of treating gastrointestinal diseases with or under endoscopy.

We encourage authors to submit their manuscripts to *WJGE*. We will give priority to manuscripts that are supported by major national and international foundations and those that are of great clinical significance.

INDEXING/ABSTRACTING

World Journal of Gastrointestinal Endoscopy is now indexed in Emerging Sources Citation Index (Web of Science), PubMed, and PubMed Central.

FLYLEAF

I-III Editorial Board

EDITORS FOR THIS ISSUE

Responsible Assistant Editor: *Xiang Li*
Responsible Electronic Editor: *Huan-Liang Wu*
Proofing Editor-in-Chief: *Lian-Sheng Ma*

Responsible Science Editor: *Jin-Xin Kong*
Proofing Editorial Office Director: *Xiu-Xia Song*

NAME OF JOURNAL
World Journal of Gastrointestinal Endoscopy

ISSN
ISSN 1948-5190 (online)

LAUNCH DATE
October 15, 2009

FREQUENCY
Monthly

EDITORS-IN-CHIEF
Atsushi Imagawa, PhD, Director, Doctor, Department of Gastroenterology, Mitoyo General Hospital, Kan-onji, Kagawa 769-1695, Japan

Juan Manuel Herrerias Gutierrez, PhD, Academic Fellow, Chief Doctor, Professor, Unidad de Gestión Clínica de Aparato Digestivo, Hospital Universitario Virgen Macarena, Sevilla 41009, Sevilla, Spain

EDITORIAL BOARD MEMBERS
All editorial board members resources online at <http://www.wjgnet.com>

www.wjgnet.com/1948-5190/editorialboard.htm

EDITORIAL OFFICE
Xiu-Xia Song, Director
World Journal of Gastrointestinal Endoscopy
Baishideng Publishing Group Inc
8226 Regency Drive, Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: editorialoffice@wjgnet.com
Help Desk: <http://www.fjgpublishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLISHER
Baishideng Publishing Group Inc
8226 Regency Drive,
Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.fjgpublishing.com/helpdesk>
<http://www.wjgnet.com>

PUBLICATION DATE
April 16, 2017

COPYRIGHT
© 2017 Baishideng Publishing Group Inc. Articles published by this Open-Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

SPECIAL STATEMENT
All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

INSTRUCTIONS TO AUTHORS
<http://www.wjgnet.com/bpg/gerinfo/204>

ONLINE SUBMISSION
<http://www.fjgpublishing.com>

Laparoscopic surgery for complex and recurrent Crohn's disease

Yusuf Sevim, Cihangir Akyol, Erman Aytac, Bilgi Baca, Orhan Bulut, Feza H Remzi

Yusuf Sevim, Department of General Surgery, Kayseri Training and Research Hospital, Kayseri 38110, Turkey

Cihangir Akyol, Division of Colorectal Surgery, Department of General Surgery, Ankara University School of Medicine, Ankara 06000, Turkey

Erman Aytac, Bilgi Baca, Department of General Surgery, Medical School, Acibadem University, Istanbul 34457, Turkey

Orhan Bulut, Department of Surgical Gastroenterology, Copenhagen University Hospital, 2650 Hvidovre, Denmark

Feza H Remzi, Inflammatory Bowel Disease Center, New York University, Langone Medical Center, New York, NY 10029, United States

Author contributions: Sevim Y, Akyol C, Aytac E, Baca B, Bulut O and Remzi FH contributed to conception and design; Sevim Y, Akyol C and Aytac E contributed to acquisition of data; Sevim Y, Akyol C, Aytac E, Baca B, Bulut O and Remzi FH contributed to analysis and interpretation of data; Sevim Y, Akyol C and Aytac E contributed to drafting the article; Sevim Y, Akyol C, Aytac E, Baca B, Bulut O and Remzi FH contributed to critical revision; Sevim Y, Akyol C, Aytac E, Baca B, Bulut O and Remzi FH approved the final version of the article to be published.

Conflict-of-interest statement: The authors have no conflict of interests.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Correspondence to: Erman Aytac, MD, FTBS, Assistant Professor of Surgery, Department of General Surgery, Medical School, Acibadem University, Büyükdere Cd. No:40 Maslak, Istanbul 34457, Turkey. ermanaytac@gmail.com

Telephone: +90-533-4144405
Fax: +90-212-2866131

Received: January 26, 2016
Peer-review started: January 27, 2016
First decision: March 23, 2016
Revised: January 3, 2017
Accepted: January 11, 2017
Article in press: January 14, 2017
Published online: April 16, 2017

Abstract

Crohn's disease (CD) is a chronic inflammatory disease of digestive tract. Approximately 70% of patients with CD require surgical intervention within 10 years of their initial diagnosis, despite advanced medical treatment alternatives including biologics, immune suppressive drugs and steroids. Refractory to medical treatment in CD patients is the common indication for surgery. Unfortunately, surgery cannot cure the disease. Minimally invasive treatment modalities can be suitable for CD patients due to the benign nature of the disease especially at the time of index surgery. However, laparoscopic management in fistulizing or recurrent disease is controversial. Intractable fibrotic strictures with obstruction, fistulas with abscess formation and hemorrhage are the surgical indications of recurrent CD, which are also complicating laparoscopic treatments. Nevertheless, laparoscopy can be performed in selected CD patients with safety, and may provide better outcomes compared to open surgery. The common complication after laparoscopic intervention is postoperative ileus seems and this may strongly relate excessive manipulation of the bowel during dissection. But additionally, unsuccessful laparoscopic attempts requiring conversion to open surgery have been a major concern due to presumed risk of worse outcomes. However, recent data show that conversions do not worsen the outcomes of colorectal surgery

in experienced hands. In conclusion, laparoscopic treatment modalities in recurrent CD patients have promising outcomes when it is used selectively.

Key words: Crohn's disease; Laparoscopic surgery; Complex disease management; Recurrent Crohn's disease

© **The Author(s) 2017.** Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: Despite advanced medical treatment alternatives including biologics, immune suppressive drugs and steroids, approximately 70% of patients with Crohn's disease (CD) require surgical intervention within 10 years of their initial diagnosis. Forty percent to 50% of patients who had an index surgery for CD require a reoperation for recurrent disease in 10 years. Index surgical treatment type and medications used after index surgery appears to be factors related to recurrence risk of CD. In experienced hands, laparoscopic approach has promising outcomes in patients with recurrent CD when it is used selectively.

Sevim Y, Akyol C, Aytac E, Baca B, Bulut O, Remzi FH. Laparoscopic surgery for complex and recurrent Crohn's disease. *World J Gastrointest Endosc* 2017; 9(4): 149-152 Available from: URL: <http://www.wjgnet.com/1948-5190/full/v9/i4/149.htm> DOI: <http://dx.doi.org/10.4253/wjge.v9.i4.149>

Crohn's disease (CD) is a chronic inflammatory disease that can develop any part of the digestive tract. CD usually arises at the terminal ileum^[1]. Despite advanced medical treatment alternatives including biologics, immune suppressive drugs and steroids, approximately 70% of patients with CD require surgical intervention within 10 years of their initial diagnosis^[2,3]. Surgery is warranted for management of medically refractory CD. Surgical treatment overcomes emergent issues, improves symptoms and patient's quality of life. Unfortunately, there is no cure for CD and it tends to recur during the disease course. Recurrent CD is described based on treatment type including medical, endoscopic or surgical.

Endoscopically documented recurrent CD can be up to 93% within one year following intestinal resection^[4], while clinically symptomatic recurrence is usually around 30% at first 3 years after surgery^[5]. Forty percent to 50% of patients who had an index surgery for CD require a reoperation for recurrent disease in 10 years^[6,7]. Index surgical treatment type and medications used after index surgery appears to be factors related to recurrence risk of CD^[2,8-10]. CD patients can be good candidates for minimally invasive treatment modalities due to the benign nature of the disease especially at the time of index surgery. However, use of laparoscopy in patients with complex CD such as extensive fistulizing or recurrent disease requiring surgical treatment is

controversial.

Majority of the surgical indications for recurrent CD are also the conditions complicating application of laparoscopic surgery such as intractable fibrotic strictures with obstruction, fistulas with abscess formation and hemorrhage^[11,12]. Based on the extension and severity of disease, surgical options including stricturoplasty, small bowel resection, ileocectomy, internal bypass, partial/total colectomy and proctectomy may be performed laparoscopically^[11,13]. In selected CD patients, laparoscopic surgery is safe, feasible and provides better outcomes compared to open surgery^[14-17]. While operative times have decreased with increased experience, operative mortality is almost none and morbidity rates ranged from 10% to 40% in patients undergoing laparoscopic surgery for recurrent CD^[17-19]. Postoperative ileus seems as the most common complication which may strongly relate excessive manipulation of the bowel during dissection^[13]. Some surgeons believe that laparoscopic approach may also provide the well-known advantages of minimally invasive surgery such as reduced postoperative pain, lower morbidity, shorter hospital stay, earlier return to daily activity, and improved quality of life in patients with recurrent CD (Table 1).

Unsuccessful laparoscopic attempts requiring conversion to open surgery have been a major concern due to presumed risk of worse outcomes and conversion rates tend to be higher in laparoscopic operations for recurrent CD^[20]. Conversion to open surgery rates varies between 6.7% and 42.3% in recurrent CD cases^[21,22]. The most common cause of conversion was adhesions^[13,23]. Having multiple resections, intraabdominal abscess and phlegmon are the other factors leading conversion in CD patients^[22]. This clinical situation raises concerns on conversion related postoperative morbidity^[24]. However, recent data show that conversions do not to worsen the outcomes of colorectal surgery in experienced hands^[25]. The data regarding to operation type and disease characteristics especially related to index resection for CD are heterogeneous in the previous reports^[26,27]. Outcomes after laparoscopic surgery for recurrent CD vary due to selection bias and experience of the surgeon^[27,28]. Laparoscopic surgery showed better outcomes with shorter length of hospital stay compared to open surgery in selected cases^[28], while laparoscopic approach did not provide expected benefits over open surgery in some series^[13,27]. Although wound complications are reduced, the benefits of laparoscopic surgery in patients with a history of previous open intestinal resection through midline laparotomy seem questionable^[13]. As an emerging technique, single incision laparoscopy can be performed for recurrent CD^[29,30]. Single incision laparoscopy can be promising in complex cases by minimizing overall wound size, decreasing unnecessary adhesiolysis for secondary port placements and it affords the surgeon the opportunity to inspect the density of adhesions through port site and lead the surgeon to convert the operation preemptively if laparoscopic surgery seems unfeasible^[31].

Table 1 Perioperative outcomes laparoscopy for complex and recurrent Crohn's disease

Ref.	Year	Patients surgery (n)	Conversion to open surgery (n)	Operative duration (min)	Hospital stay (d)
Wu <i>et al</i> ^[17]	1997	CL: 14	1	152	4.8
		RL: 10	2	144	3.9
		PL: 22	2	139	4.5
		O: 70	(-)	202 ^a	7.9 ^a
Hasegawa <i>et al</i> ^[28]	2003	RL: 16	2	210 ^a	6.0
		PL: 45	3	180	8.0
Moorthy <i>et al</i> ^[22]	2004	RL: 26	11	118	8.0
		PL: 31	4	127	7.0
Goyer <i>et al</i> ^[32]	2009	Comp: 54 ^b	20 ^a	214 ^a	8.0
		Uncomp: 70	10	191	7.0
Chaudhary <i>et al</i> ^[21]	2010	RL: 30	2	125 ^a	3.0
		PL: 29	3	85	3.0
Brouquet <i>et al</i> ^[27]	2010	L: 29	9	215	9.0
		O: 33		226	9.0
Pinto <i>et al</i> ^[18]	2011	RL: 50	16	201	7.4
		PL: 80	15	182	6.7
Aytac <i>et al</i> ^[13]	2012	L: 26	3	169	6.4
		O: 26		158	6.9
Huang <i>et al</i> ^[20]	2012	RL: 48	10	100	ND
		PL: 82	14	106	ND

^aBold: Statistically significant; ^b27 of these patients had recurrent disease. CL: Laparoscopic surgery for complicated disease (phlegmon, abscess); PL: Laparoscopic surgery for primary disease; RL: Laparoscopic surgery for recurrent disease; L: Laparoscopic surgery; O: Open surgery; Comp: Complicated; Uncomp: Uncomplicated.

In experienced hands, laparoscopic approach has promising outcomes in patients with recurrent CD when it is used selectively. There is a need for new studies which focus on identification of proper patients who may benefit from laparoscopic surgery for recurrent and complex CD.

REFERENCES

- Longobardi T, Jacobs P, Bernstein CN. Work losses related to inflammatory bowel disease in the United States: results from the National Health Interview Survey. *Am J Gastroenterol* 2003; **98**: 1064-1072 [PMID: 12809829 DOI: 10.1111/j.1572-0241.2003.07285.x]
- Bernell O, Lapidus A, Hellers G. Recurrence after colectomy in Crohn's colitis. *Dis Colon Rectum* 2001; **44**: 647-654; discussion 654 [PMID: 11357022]
- Vind I, Riis L, Jess T, Knudsen E, Pedersen N, Elkjaer M, Bak Andersen I, Wewer V, Nørregaard P, Moesgaard F, Bendtsen F, Munkholm P. Increasing incidences of inflammatory bowel disease and decreasing surgery rates in Copenhagen City and County, 2003-2005: a population-based study from the Danish Crohn colitis database. *Am J Gastroenterol* 2006; **101**: 1274-1282 [PMID: 16771949 DOI: 10.1111/j.1572-0241.2006.00552.x]
- Olaion G, Smedh K, Sjö Dahl R. Natural course of Crohn's disease after ileocolic resection: endoscopically visualised ileal ulcers preceding symptoms. *Gut* 1992; **33**: 331-335 [PMID: 1568651]
- Sachar DB. The problem of postoperative recurrence of Crohn's disease. *Med Clin North Am* 1990; **74**: 183-188 [PMID: 2404175]
- Nordgren SR, Fasth SB, Oresland TO, Hultén LA. Long-term follow-up in Crohn's disease. Mortality, morbidity, and functional status. *Scand J Gastroenterol* 1994; **29**: 1122-1128 [PMID: 7886401]
- Aratari A, Papi C, Leandro G, Viscido A, Capurso L, Caprilli R. Early versus late surgery for ileo-caecal Crohn's disease. *Aliment Pharmacol Ther* 2007; **26**: 1303-1312 [PMID: 17848181 DOI: 10.1111/j.1365-2036.2007.03515.x]
- Renna S, Cammà C, Modesto I, Cabibbo G, Scimeca D, Civitavecchia G, Moccia F, Orlando A, Enea M, Cottone M. Meta-analysis of the placebo rates of clinical relapse and severe endoscopic recurrence in postoperative Crohn's disease. *Gastroenterology* 2008; **135**: 1500-1509 [PMID: 18823987 DOI: 10.1053/j.gastro.2008.07.066]
- Borley NR, Mortensen NJ, Chaudry MA, Mohammed S, Warren BF, George BD, Clark T, Jewell DP, Kettlewell MG. Recurrence after abdominal surgery for Crohn's disease: relationship to disease site and surgical procedure. *Dis Colon Rectum* 2002; **45**: 377-383 [PMID: 12068198]
- Rutgeerts P, Goobes K, Peeters M, Hiele M, Penninckx F, Aerts R, Kerremans R, Vantrappen G. Effect of faecal stream diversion on recurrence of Crohn's disease in the neoterminal ileum. *Lancet* 1991; **338**: 771-774 [PMID: 1681159]
- Jones DW, Finlayson SR. Trends in surgery for Crohn's disease in the era of infliximab. *Ann Surg* 2010; **252**: 307-312 [PMID: 20585239 DOI: 10.1097/SLA.0b013e3181e61df5]
- Strong SA, Koltun WA, Hyman NH, Buie WD. Practice parameters for the surgical management of Crohn's disease. *Dis Colon Rectum* 2007; **50**: 1735-1746 [PMID: 17690937 DOI: 10.1007/s10350-007-9012-7]
- Aytac E, Stocchi L, Remzi FH, Kiran RP. Is laparoscopic surgery for recurrent Crohn's disease beneficial in patients with previous primary resection through midline laparotomy? A case-matched study. *Surg Endosc* 2012; **26**: 3552-3556 [PMID: 22648125 DOI: 10.1007/s00464-012-2361-x]
- Alabaz O, Iroatulam AJ, Nessim A, Weiss EG, Nogueras JJ, Wexner SD. Comparison of laparoscopically assisted and conventional ileocolic resection for Crohn's disease. *Eur J Surg* 2000; **166**: 213-217 [PMID: 10755335 DOI: 10.1080/110241500750009302]
- Bemelman WA, Slors JF, Dunker MS, van Hogeand RA, van Deventer SJ, Ringers J, Griffioen G, Gouma DJ. Laparoscopic-assisted vs. open ileocolic resection for Crohn's disease. A comparative study. *Surg Endosc* 2000; **14**: 721-725 [PMID: 10954817]
- Schmidt CM, Talamini MA, Kaufman HS, Lilliemoe KD, Learn P, Bayless T. Laparoscopic surgery for Crohn's disease: reasons for conversion. *Ann Surg* 2001; **233**: 733-739 [PMID: 11371731]
- Wu JS, Birnbaum EH, Kodner IJ, Fry RD, Read TE, Fleshman JW. Laparoscopic-assisted ileocolic resections in patients with Crohn's disease: are abscesses, phlegmons, or recurrent disease contraindications? *Surgery* 1997; **122**: 682-688; discussion 682-688 [PMID: 9347843]
- Pinto RA, Shawki S, Narita K, Weiss EG, Wexner SD. Laparoscopy for recurrent Crohn's disease: how do the results compare with the results for primary Crohn's disease? *Colorectal Dis* 2011; **13**: 302-307 [PMID: 19912288 DOI: 10.1111/j.1463-1318.2009.02133.x]
- Tavernier M, Lebreton G, Alves A. Laparoscopic surgery for complex Crohn's disease. *J Visc Surg* 2013; **150**: 389-393 [PMID: 24119432 DOI: 10.1016/j.jvisc.2013.09.004]
- Huang R, Valerian BT, Lee EC. Laparoscopic approach in patients with recurrent Crohn's disease. *Am Surg* 2012; **78**: 595-599 [PMID: 22546134]
- Chaudhary B, Glancy D, Dixon AR. Laparoscopic surgery for recurrent ileocolic Crohn's disease is as safe and effective as primary resection. *Colorectal Dis* 2011; **13**: 1413-1416 [PMID: 21087388 DOI: 10.1111/j.1463-1318.2010.02511.x]
- Moorthy K, Shaul T, Foley RJ. Factors that predict conversion in patients undergoing laparoscopic surgery for Crohn's disease. *Am J Surg* 2004; **187**: 47-51 [PMID: 14706585]
- Holubar SD, Dozois EJ, Privitera A, Cima RR, Pemberton JH, Young-Fadok T, Larson DW. Laparoscopic surgery for recurrent ileocolic Crohn's disease. *Inflamm Bowel Dis* 2010; **16**: 1382-1386 [PMID: 20027655 DOI: 10.1002/ibd.21186]
- Gonzalez R, Smith CD, Mason E, Duncan T, Wilson R, Miller J, Ramshaw BJ. Consequences of conversion in laparoscopic colorectal surgery. *Dis Colon Rectum* 2006; **49**: 197-204 [PMID: 1681159]

- 16328607 DOI: 10.1007/s10350-005-0258-7]
- 25 **Aytac E**, Stocchi L, Ozdemir Y, Kiran RP. Factors affecting morbidity after conversion of laparoscopic colorectal resections. *Br J Surg* 2013; **100**: 1641-1648
 - 26 **Bandyopadhyay D**, Sagar PM, Mirnezami A, Lengyel J, Morrison C, Gatt M. Laparoscopic resection for recurrent Crohn's disease: safety, feasibility and short-term outcomes. *Colorectal Dis* 2011; **13**: 161-165 [PMID: 19888954 DOI: 10.1111/j.1463-1318.2009.02100.x]
 - 27 **Brouquet A**, Bretagnol F, Soprani A, Valleur P, Bouhnik Y, Panis Y. A laparoscopic approach to iterative ileocolonic resection for the recurrence of Crohn's disease. *Surg Endosc* 2010; **24**: 879-887 [PMID: 19730944 DOI: 10.1007/s00464-009-0682-1]
 - 28 **Hasegawa H**, Watanabe M, Nishibori H, Okabayashi K, Hibi T, Kitajima M. Laparoscopic surgery for recurrent Crohn's disease. *Br J Surg* 2003; **90**: 970-973 [PMID: 12905550 DOI: 10.1002/bjs.4136]
 - 29 **Kawahara H**, Watanabe K, Ushigome T, Noaki R, Kobayashi S, Yanaga K. Single-incision laparoscopic right colectomy for recurrent Crohn's disease. *Hepatogastroenterology* 2010; **57**: 1170-1172 [PMID: 21410052]
 - 30 **Moftah M**, Nazour F, Cunningham M, Cahill RA. Single port laparoscopic surgery for patients with complex and recurrent Crohn's disease. *J Crohns Colitis* 2014; **8**: 1055-1061 [PMID: 24589026 DOI: 10.1016/j.crohns.2014.02.003]
 - 31 **Aytac E**, Stocchi L, Williams R, Remzi FH, Costedio MM. Single-port laparoscopic fecal diversion: more than cosmetic benefits? *Surg Laparosc Endosc Percutan Tech* 2014; **24**: e133-136
 - 32 **Goyer P**, Alves A, Bretagnol F, Bouhnik Y, Valleur P, Panis Y. Impact of complex Crohn's disease on the outcome of laparoscopic ileocecal resection: a comparative clinical study in 124 patients. *Dis Colon Rectum* 2009; **52**: 205-210 [PMID: 19279413 DOI: 10.1007/DCR.0b013e31819c9c08]

P- Reviewer: Saniabadi AR, Vradelis S **S- Editor:** Song XX
L- Editor: A **E- Editor:** Wu HL





Published by **Baishideng Publishing Group Inc**

8226 Regency Drive, Pleasanton, CA 94588, USA

Telephone: +1-925-223-8242

Fax: +1-925-223-8243

E-mail: bpgoffice@wjgnet.com

Help Desk: <http://www.f6publishing.com/helpdesk>

<http://www.f6publishing.com>

